

## PR-9

**INVESTIGATION ON SYNTHESIS, STRUCTURAL  
AND OPTICAL PROPERTIES OF CDS NANOPARTICLES**

**G. Thirumala Rao,<sup>1\*</sup> R. V. S. S. N. Ravikumar,<sup>2</sup> N. Bakthavatchala Reddy,<sup>3</sup>  
Grigory V Zyryanov<sup>3,4</sup>**

<sup>1</sup>*Physics division, Department of Basic Science & Humanities,  
GMR Institute of Technology, GMR Nagar, Rajam-532 127, A.P., India.*  
<sup>2</sup>*Department of Physics, Acharya Nagarjuna University, Nagarjuna Nagar,  
Guntur-522510, A.P., India.*

<sup>3</sup>*Ural Federal University, Chemical Engineering Institute Yekaterinburg,  
620002, Russian Federation.*

<sup>4</sup>*I. Ya. Postovskiy Institute of Organic Synthesis,  
Ural Division of the Russian Academy of Sciences, 22 S. Kovalevskoy St.,  
620219 Yekaterinburg, Russian Federation.*

\*Corresponding author e-mail: thirumalarao.g@gmr.it.org, thirumalaphy@gmail.com

**Abstract.** Hexagonal wurtzite structured CdS nanoparticles were synthesized by using chemical precipitation method. Various spectroscopic technics were used for the characterization of the prepared sample. X-ray diffraction pattern shows the high intense crystalline peaks and the average crystallite size is found to be 19 nm. FT-IR spectrum exhibited a metal sulfide band at  $615\text{ cm}^{-1}$  and other functional groups. SEM micrographs reveal the non-uniformly distributed spherical shaped structures. EDS analysis confirms the stoichiometric composition and presence of target elements. Optical absorption spectrum exhibited a broad peak at 510 nm, which indicates the shifting of absorption range of CdS nanoparticles into the visible region. PL spectrum exhibited the characteristic emission bands in visible region.